AN IMPROVED METHOD FOR ISOLATION OF *THIELAVIOPSIS PARADOXA* FROM STEM BLEEDING AFFECTED COCONUT PALMS

Anil Kumar and K.K.N. Nambiar¹

Stem bleeding of coconut caused by *Thielaviopsis paradoxa* (de Seynes) Von Hohnel is an important disease affecting coconut in many countries (Menon and Pandalai, 1958; Ohler, 1966, Nambiar and Sastry, 1988). Isolation of the pathogen from diseased tissues on different agar based media has given inconsistent results (Anon., 1976; Anon., 1986), None of the selective media reported for certain related fungi viz., *T. basicola* (Berk and Br.) Ferr. (David, 1978;Tsao and Bricker, 1966) and *Ceratocystis wagnerii* Goheen and Cobb (Hicks et al., *1980) proved* useful for the isolation of *T. paradoxa* Since, standardization of an isolation method is a basic necessity for any study on plant pathogens, an attempt was made to improve upon existing methods.

MATERIALS AND METHODS

Samples were collected from. 25 diseased palms near CPCRI, Kasaragod. Small pieces (0.25 x 0.25 cm), cut from margin of infected stem tissues were sterilized with 0.1 % HgC12 for 30 secs and washed thoroughly with sterile water. Two media viz. potato dextrose agar (PDA) and sugarcane juice agar (SJA; 200 ml sugarcanc juice, 20 g agar powder and 800 ml distilled water; pH 4.5), with and without addition of antimicrobial agents were compared with a bait method for the isolation of *T. paradoxa* Certain antimicrobial agents viz., sodium-diethyldithiocarbamate (100 ppm), sodium propionate (1,000 ppm), oxbile (1,000 ppm), streptomycin (100 ppm), tetracycline (50 ppm) and penicillin (100 ppm) were added to media after screening a number of chemicals for their efficacy (Table 1). Infected bits were kept on agar surface of plates containing different media. Frond pieces (10-12 cm in length) from old coconut leaves were found most effective as bait (Table 2) and were used in subsequent studies. The pieces were inoculated with infected bits by bore-hole method (Nambiar *et al.*, 1985), incubated at 30'C in polythene bags and isolations were made after 10 days of incubation from margin of rotten frond tissues on SJA. Response of *T. parado* to different media and baiting method was recorded.

RESULTS AND DISCUSSION

The data on successful isolation of *T. paradoxa* by different methods are presented in Table 3. The pathogen was isolated from 98.3% infected bits by bait method in comparison to 8.0-19.6% isolation on various media tested. SJA proved slightly better medium over PDA for the purpose. Addition of selected antimicrobial agents to the media did not favour the selective isolation of *T. paradoxa*_Colonies of many fungi viz., *Aspergillu* sp., *Trichoderma* sp., *Pestalotia* sp., Phoma sp. etc. were noticed in isolation plates of all agar-based media. Such contamination problem could be avoided by using bait-method. The failure of selective media, reported for isolation of related fungi, for obtaining isolation of *T. paradox* can be explained on the basis of observations presented in Table 1. Cycloheximide, Brassicol (pentachloro-nitrobenzene) and mycostatin at 100 ppm and Rose Bengal at

50 ppm inhibited chlarnydospore germination or mycelial growth of T. paradox or both to appreciable levels. These chernicals have been used in the selective media for T. basicola (David, 1978, Tsao, and Bricker, 1966) and C. wagnerii (Hicks 1980) at much higher contrations.

¹ Scientists, N.R.C.A.F., Pahug Dam, Jhansi (UP) India and Central Plantation Crops Research Institute, Kasaragod, India, respectively.

	Chlamydospore germination/ mycelial growth at					
Chemicals	10 ppm		50 ppm		100 ppm	
	Germination	Growth*	Germination	Growth	Germination	Growth
Antifungal compouds						
Bavistin 50 WP	87.00	0.00	86.4	0.0	81.8	0.0
Brassicol 75 WP	15.2	52.5	17.6	45.0	8.9	40.0
Captaf 75 WP	98.6	0.0	94.1	0.0	91.5	0.0
Copper sulphate	100.0	100.0	63.5	91.3	55.0	57.5
Dithane M-45 (75%)	17.8	65.2	0.0	55.1	0.0	50.6
Foltaf 80 WP	0.0	35.0	0.0	28.8	0.0	27.5
Murcuric chloride	17.4	73.0	0.0	37.1	0.0	8.9
Ox-bile	99.1	100.0	86.7	100.0	87.6	97.2
Sodium-diethyl-dith- iocarbamate (Technical)	100.0	69.6 98.6	100.0	67.4 97 3	79.6 94 5	65.1 89.6
Vitavax 75 WP	23.6	73.7	0.0	30.0	0.0	16.3
Antibiotics	25.0	15.1	0.0	50.0	0.0	10.5
Aureofungin Sol (30%)	33 /	563	0.0	0.0	0.0	0.0
Cyclobevimide	0.0	37.5	0.0	16.8	0.0	6.8
Mycostatin (4,960- units/mg)	16.8	34.8	0.0	16.8	0.0	6.8
salt > 1435 units/mg) Pimaricin (2.5 aqueous-	100.0	96.3	99.5	92.5	96.4	80.0
susp.)	51.2	31.4	29.7	30.0	0.0	26.6
Streptomycin sulphate	97.1	97.2	98.2	97.2	100.0	97.2
hydrochloride	90.4	100.0	87.5	100.0	92.7	100.0
Vancomycin	90.1	75.0	81.3	76.6	79.6	72.4
Organic dyes						
Rose bengal	99.3	25.8	0.0	24.7	0.0	24.5
Melachite green	0.0	21.3	0.0	17.5	0.0	17.5

Table I. Effect of some antifungal compounds, antibiotics and organic dyes on chlamydospore germination and mycelial growth of *T. paradoxa*

*Expressed as percentage of growth recorded on control medium @ Average of three replications

artificial moculation with 1. puralox							
Leaf position	Extent of colonization* after						
	4 days	8 days	12 days	16 days			
2 (upper whorl)	-	+	+++	++++			
12 (middle whort)	-	-	+	++			
26 (lower whorl)	+	+++	++++	++++			

Table 2: Effect of age on extent of coconut leaf frond fissue colonization after artificial inoculation with T. paradox

*-: No decay; + : Decay up to 25%; ++ : Decay from 26 to 50%; +++: Decay from 51 to 75%; ++++ : Decay 75%

Table 3: Details of isolation of T. paradox from stem bleeding affected coconut palms by using different media/ method

Media/method	No of bits from which <i>T. paradoxa</i> was isolated	No. of palms from which <i>T</i> . <i>paradox</i> was isolated		
PDA	21* (8.4)#	11(44)		
Amencled PDA	20(8.0)	13(52)		
SJA	49(19.6)	22(88)		
Amencled SJA	28(10.8)	17(68)		
Host bait method	248(99.2)	25(100)		

* 250 bits from 25 diseased palms (10 bits/palm) were used for isolation.

Figure in the parenthesis inclicate the percent isolation.

CONCLUSIONS

- 1. The results clearly bring out the advantage of using bait method for isolation of *T. paradoxa* from stern bleeding affected palms.
- 2. The method is highly reproduciable and by using it, continous association of *T. paradox* with the diseased palms could be established (Table 3). In past, the inconsistent isolations had created a lot of confusion regarding etiology of the disease (Anon., 1976; Anon., 1986). Thus, this confirms the finding of Nambiar *et al.* (1985), who successfully reproduced the disease symptoms by inoculating healthy trees with *T. paradoxa*

ACKNOWLEDGEMENTS

The authors are thankful to Dr. M.K. Nair, the Director. CPCRI, Kasaragod for providing encouragement and facilities.

REFERENCES

- 1. ANON, 1976. Stern bleeding disease of *coconut. CPCRI Annual Report* CPCRI, Kasaragod, p. 62.
- 2. ANON, 1986. Stern bleeding disease of coconut. CPCRI *Annual Report* CPCRI, Kasaragod, p. 19.
- 3. DAVID, C.H. HSI. 1978. Effect of crop sequence, previous peanut blackhull severity, and time of sampling on soil population of *Thielaviopsis basicola*. *Phypatholog* 68: 1442-1445.
- 4. HICKS, B.R., COBB, F.W.R., AND GERSPER. P.L. 1980. Isolation of *Ceratocystis wagener* from forest soil with a selective m.edium. *Phytopathology* 70: 880-883.
- 5. MENON, K.P.V., and PANDALAI, K.M. 1960. The coconut palm A monograph. Published by Indian Central Coconut Committee p. 384.
- 6. NAMBIAR, K.K.N., JOSHI, Y., VENUGOPAL, M.N., and MOHAN, R.C. 1985. Stem bleeding disease of coconut: Reproduction of symptoms by inoculation with *Thielaviopsis paradoxa*. J. *Plantation Crop* 14: 130-133.
- 7. NAMBIAR, K.K.N., and SASTRY, R.K. 1988. Stern bleeding disease of coconut: Current status and approaches for its control. *Philippine J. Coconut Studies* 13(1) : 30-32.
- 8. OHLER, J.G. 1964. Coconut-Tree of Life. Published by FAO. Rome. p. 446.
- 9. TSAO, P.H., and BRICKER, J.L. 1966. Chlamyclospores of *Thielaviopsis* basicola as surviving propagules in natural soils. Phytopathology 56(9): 1012-1014.